

Table 8—Critical Values for the Sign Test

Reject the null hypothesis when the test statistic x is less than or equal to the value in the table.

<i>n</i>	One-tailed, $\alpha = 0.005$		$\alpha = 0.01$	$\alpha = 0.025$	$\alpha = 0.05$
	Two-tailed, $\alpha = 0.01$	$\alpha = 0.02$	$\alpha = 0.05$	$\alpha = 0.10$	
8	0	0	0	1	
9	0	0	1	1	
10	0	0	1	1	
11	0	1	1	2	
12	1	1	2	2	
13	1	1	2	3	
14	1	2	3	3	
15	2	2	3	3	
16	2	2	3	4	
17	2	3	4	4	
18	3	3	4	5	
19	3	4	4	5	
20	3	4	5	5	
21	4	4	5	6	
22	4	5	5	6	
23	4	5	6	7	
24	5	5	6	7	
25	5	6	6	7	

Note: Table 8 is for one-tailed or two-tailed tests.

The sample size n represents the total number of + and - signs. The test value is the smaller number of + or - signs.

From *Journal of American Statistical Association* Vol. 41 (1946), pp. 557–566. W. J. Dixon and A. M. Mood.

Table 9—Critical Values for the Wilcoxon Signed-Rank Test

Reject the null hypothesis when the test statistic w_s is less than or equal to the value in the table.

<i>n</i>	One-tailed, $\alpha = 0.05$		$\alpha = 0.025$	$\alpha = 0.01$	$\alpha = 0.005$
	Two-tailed, $\alpha = 0.10$	$\alpha = 0.05$	$\alpha = 0.02$	$\alpha = 0.01$	
5	1	—	—	—	—
6	2	1	—	—	—
7	4	2	0	—	—
8	6	4	2	0	
9	8	6	3	2	
10	11	8	5	3	
11	14	11	7	5	
12	17	14	10	7	
13	21	17	13	10	
14	26	21	16	13	
15	30	25	20	16	
16	36	30	24	19	
17	41	35	28	23	
18	47	40	33	28	
19	54	46	38	32	
20	60	52	43	37	
21	68	59	49	43	
22	75	66	56	49	
23	83	73	62	55	
24	92	81	69	61	
25	101	90	77	68	
26	110	98	85	76	
27	120	107	93	84	
28	130	117	102	92	
29	141	127	111	100	
30	152	137	120	109	

From *Some Rapid Approximate Statistical Procedures*. Copyright 1949, 1964 Lederle Laboratories, American Cyanamid Co., Wayne, N.J.